arXiv:hep-th/0305003v3 11 Aug 2003

Noncommutative massive Thirring model in three-dimensional spacetime

T. Mariz, J.R. Nascimento and R.F. Ribeiro
Departamento de Física, Universidade Federal da Paraíba,
Caixa Postal 5008, 58051-970 João Pessoa, Paraíba, Brazil

F.A. Brito
Departamento de Ciências Físicas e Biológicas,
Universidade Regional do Cariri, 63100-000 Crato, Ceará, Brazil

(Dated: February 1, 2008)

Abstract

We evaluate the noncommutative Chern-Simons action induced by fermions interacting with an Abelian gauge field in a noncommutative massive Thirring model in (2+1)-dimensional spacetime. This calculation is performed in the Dirac and Majorana representations. We observe that in Majorana representation when $\bar{\theta}$ goes to zero we do not have induced Chern-Simons term in the dimensional regularization scheme.

Noncommutative field theories have been intensively studied during recent years — for a review in noncommutative field theory see, for instance, [1, 2] and references therein. At one hand, noncommutative spacetime, and in turn noncommutative field theory, arise as a particular low energy limit of string theory [3, 4, 5]. On the other hand, in condensed matter theory, there is an example leading directly to the noncommutative geometry. This is the example of the theory of the electrons in a constant magnetic field, projected to the lowest Landau level, which is naturally thought of as a noncommutative field theory [6, 7]. The noncommutativity geometry plays a very important role in the context of the Hall effect [8, 9].

The noncommutativity of the spacetime can be postulated by the following commutation relation

$$[x^\mu, x^\nu] \;=\; i\theta^{\mu\nu} \tag{1}$$

with a parameter $\theta^{\mu\nu}$ which is a constant antisymmetric tensor, and it has the canonical dimension of inverse mass squared. As a consequence of the noncommutativity of the spacetime coordinates one replaces the ordinary product of functions by the Moyal product

$$f(x) \star g(x) \;=\; e^{\frac{i}{2}\theta^{\mu\nu}\partial_\mu \partial'_\nu} f(x) g(x')|_{x'=x} \tag{2}$$

which is associative, noncommutative and satisfies the following relation

$$\int f(x) \star g(x) = \int g(x) \star f(x) = \int fg \tag{3}$$

that is a consequence of the momentum conservation. Thus to formulate a field theory on a noncommutative spacetime it simply needs to replace the usual multiplication function by the Moyal product.

In this works we are interested in analyzing the induction of the Chern-Simons action in (2+1)-dimensional spacetime for the noncommutative massive Thirring model. There is a well known class of (2+1)-dimensional spacetime theories exhibiting interesting phenomena such as exotic statistics, fractional spin and massive gauge fields [10, 11]. All these phenomena are of topological nature and they can be produced when we add Chern-Simons term to the Lagrangian which describe the system under consideration. Another important point that has been analyzed by a great number of authors is to understand the mechanism by which the Chern-Simons term can be generated in the usual spacetime. Particularly, the induction of the Chern-Simons term via radiative correction has been explicitly verified in the context of models with four-fermion (like Gross-Neveu) [12] or current-current (like Thirring) [13] self-interaction in the commutative spacetime. The consideration of such a class of three-dimensional theories in noncommutative spacetime may reveal new interesting physics.

We consider two different representations for the covariant derivative of the fermions, namely, the Dirac and Majorana representations. We discuss some relevant properties of the induced noncommutative Chern-Simons actions [14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24]. In the Majorana representation we shall discuss the Chern-Simons action dependence on the noncommutative parameter $\theta_{\mu\nu}$. Our investigations in this paper follows the lines of the papers [25, 26, 27, 28] which make use of the derivative expansion method. In order to perform the divergent integrals we apply the dimensional regularization.

The action for noncommutative massive Thirring model takes the form

$$S \;=\; \int \mathrm{d}^3x \left[\bar{\psi} \star (i\partial\!\!\!/ - m)\psi - \frac{g}{2N}(\bar{\psi}\gamma_\mu \star \psi) \star (\bar{\psi}\gamma^\mu \star \psi)\right]. \tag{4}$$

The most efficient way to implement the $1/N$ expansion for this model is to introduce an auxiliary field, $A_\mu = (g/\sqrt{N})\bar{\psi}\gamma_\mu\star\psi$, in order to eliminate the current-current self-interaction in Eq.(4) so that it becomes

$$S[A,m] \;=\; \int \mathrm{d}^3x \left[\bar{\psi} \star (i\mathcal{D}\!\!\!\!/ - m)\psi + \frac{1}{2g}A_\mu \star A^\mu\right], \tag{5}$$

where the covariant derivative acting on ψ is given by

$$\mathcal{D}_\mu\psi = \begin{cases} \partial_\mu\psi + ig_{\mathrm{D}}\psi \star A_\mu, & \text{Dirac representation,} \\ \partial_\mu\psi - ig_{\mathrm{M}}[A_\mu, \psi]_\star, & \text{Majorana representation,} \end{cases} \tag{6}$$

with $g_{\mathrm{D}} = \frac{1}{\sqrt{N}}$ and $g_{\mathrm{M}} = \frac{1}{2\sqrt{N}}$. The γ-matrices in (2+1) dimensions are Pauli matrices which satisfy the well known algebra

$$\gamma^\mu\gamma^\nu = g^{\mu\nu} - i\epsilon^{\mu\nu\lambda}\gamma_\lambda \tag{7}$$

where $g^{\mu\nu} = (+,-,-)$ and $\epsilon^{123} = +1$.

The effective action $\Gamma(A,m)$ is defined as

$$Z(A,m) = e^{i\Gamma(A,m)} = \int \mathcal{D}\psi\mathcal{D}\bar{\psi}e^{iS[A,m]}. \tag{8}$$

We shall calculate the effective action integrating on the fermions in the Dirac representation. Let us make use of the derivative expansion method to evaluate the one fermion-loop effective action. To do this we substitute the Eq.(5) into Eq.(8), and use the Moyal product, Eq.(2). Thus, the effective action in the momentum space is given by

$$\Gamma[A,m] = -iN\,\mathrm{Tr}\,\ln\Big(\not{p} - m - g_{\mathrm{D}}\; e^{-\partial\times p}\,\not{A}(x)\Big). \tag{9}$$

The Tr stands for the trace over Dirac matrices as well as trace over the integration in momentum and coordinate spaces. $A_\mu(x)$ being dependent on the position do not commute with functions of momentum and it is not clear how to separate out the momentum and space dependent quantities. To do that, we shall use the techniques of derivative expansion [25, 26, 27, 28] that proceeds as follows:

$$\Gamma[A,m] = \Gamma[m] + \Gamma'[A,m]$$

where the first term is $\Gamma[m] = -iN\,\mathrm{Tr}\,\ln(\not{p} - m)$, which does not depend on the auxiliary field A. We concentrate ourselves on the second term, which is known as Matthews-Salam determinant [29]. The term is given by

$$\Gamma'[A,m] = -iN\,\mathrm{Tr}\,\ln\left(1 - g_{\mathrm{D}}\frac{1}{\not{p}-m}\; e^{-\partial\times p}\,\not{A}(x)\right). \tag{10}$$

The new effective action can be written as

$$\Gamma'[A,m] \;=\; iN\,\mathrm{Tr}\sum_{n=1}^{\infty}\frac{1}{n}\Big[g_{\mathrm{D}}\,S(p)\;e^{-\partial\times p}\,\not{A}(x)\Big]^n \tag{11}$$

where $S(p) = (\not{p} - m)^{-1}$ is the free fermion propagator. The exponential term that appears in the Eq.(11) is due to the noncommutativity.

The corresponding action for the quadratic term in A_μ is given by

$$\Gamma^{(2)}[A,m] \;=\; \frac{i}{2}\,\mathrm{Tr}\frac{1}{\not{p}-m}\,e^{-\partial\times p}\,\not{A}(x)\frac{1}{\not{p}-m}\,e^{-\partial'\times p}\,\not{A}(x')\Big|_{x'=x}. \tag{12}$$

Let us use the following identity in order to disentangle the x and p trace, i.e.,

$$\not{A}S(p)e^{-\partial'\times p} = \gamma^\mu S(p-i\partial)e^{-\partial'\times(p-i\partial)}A_\mu. \tag{13}$$

Now we can rewrite the Eq.(12) in the form

$$\Gamma^{(2)} \;=\; \frac{i}{2}\int \mathrm{d}^3x\,\Pi^{\mu\nu}A_\nu(x')\star A_\mu(x)\Big|_{x'=x} \tag{14}$$

where the tensor $\Pi^{\mu\nu}$ is given by

$$\Pi^{\mu\nu} \;=\; \mathrm{tr}\int\frac{\mathrm{d}^3p}{(2\pi)^3}S(p)\gamma^\mu S(p-i\partial)\gamma^\nu e^{-(\partial+\partial')\times p}. \tag{15}$$

Expanding $S(p-i\partial)$ around p,

$$S(p-i\partial) \;=\; S(p)+S(p)i\not{\partial}S(p)+\cdots \tag{16}$$

and keeping only terms up to the first order in the derivative ∂, the Eq.(15) is written as

$$\Pi^{\mu\nu} \;=\; \mathrm{tr}\int\frac{\mathrm{d}^3p}{(2\pi)^3}S(p)\gamma^\mu S(p)i\not{\partial}S(p)\gamma^\nu. \tag{17}$$

By manipulating this equation is straightforward to get

$$\Pi^{\mu\nu} \;=\; im\,\mathrm{tr}\int\frac{\mathrm{d}^3p}{(2\pi)^3}\frac{1}{(p^2-m^2)^3}\times\left(\not{p}\gamma^\mu\not{p}\not{\partial}\gamma^\nu+\not{p}\gamma^\mu\not{\partial}\not{p}\gamma^\nu+\gamma^\mu\not{p}\not{\partial}\not{p}\gamma^\nu+m^2\gamma^\mu\not{\partial}\gamma^\nu\right). \tag{18}$$

Now applying the tr on the γ matrices and solving the momentum integration, which is finite, we get to the result

$$\Gamma^{(2)} \;=\; -\frac{1}{8\pi}\frac{m}{|m|}\int\mathrm{d}^3x\epsilon^{\mu\nu\rho}\partial_\mu A_\nu\star A_\rho. \tag{19}$$

The corresponding action for the cubic term in A_μ is given by

$$\Gamma^{(3)}[A,m] \;=\; \frac{ig_{\mathrm{D}}}{3}\,\mathrm{Tr}\,\frac{1}{\not{p}-m}\,e^{-\partial\times p}\,\not{A}(x)\,\frac{1}{\not{p}-m}\,e^{-\partial'\times p}\,\not{A}(x')\,\frac{1}{\not{p}-m}\,e^{-\partial''\times p}\,\not{A}(x'')\Big|_{x'=x''=x}. \tag{20}$$

Using the equations (13) and (16) we can write the Eq.(20) as

$$\Gamma^{(3)} \;=\; \frac{ig_{\mathrm{D}}}{3}\int\mathrm{d}^3x\,\Gamma^{\mu\rho\nu}A_\nu\star A_\rho\star A_\mu \tag{21}$$

where the tensor $\Gamma^{\mu\rho\nu}$ is

$$\Gamma^{\mu\rho\nu} \;=\; \mathrm{tr}\int\frac{\mathrm{d}^3p}{(2\pi)^3}S(p)\gamma^\mu S(p)\gamma^\rho S(p)\gamma^\nu, \tag{22}$$

Now applying the tr on the γ matrices and calculating the momentum integration in the Eq.(22) we obtain

$$\Gamma^{(3)} \;=\; \frac{ig_{\mathrm{D}}}{12\pi}\frac{m}{|m|}\int\mathrm{d}^3x\epsilon^{\mu\nu\rho}A_\mu\star A_\nu\star A_\rho. \tag{23}$$

Finally, combining both contributions, Eq.(19) and Eq.(23), we find the result

$$\Gamma_{\mathrm{cs}}[A] \;=\; -\frac{1}{8\pi}\frac{m}{|m|}\int\mathrm{d}^3x\epsilon^{\mu\nu\rho}\left(\partial_\mu A_\nu\star A_\rho-\frac{2ig_{\mathrm{D}}}{3}A_\mu\star A_\nu\star A_\rho\right), \tag{24}$$

which is the Chern-Simons action for the Dirac representation. Observe that even in the abelian case we have a cubic term, similar to the Chern-Simons action for ordinary non-abelian case.

In the Majorana representation the effective action takes the form

$$\tilde{\Gamma}'[A,m] = i\frac{N}{2}\mathrm{Tr}\sum_{n=1}^{\infty}\frac{1}{n}\left[g_{\mathrm{M}}\,S(p)\,\left(e^{-\partial\times p}-e^{\partial\times p}\right)\not{A}(x)\right]^{n}. \tag{25}$$

The corresponding action for the quadratic term in A is given by

$$\begin{aligned}\tilde{\Gamma}^{(2)} &= \frac{i}{4}\frac{g_{\mathrm{M}}^2}{g_{\mathrm{D}}^2}\int \mathrm{d}^3x\,\mathrm{tr}\int\frac{\mathrm{d}^3p}{(2\pi)^3}S(p)\gamma^{\mu}S(p-i\partial)\gamma^{\nu}\\ &\times\ \left[\left(e^{(\partial+\partial')\times p}-e^{-(\partial-\partial')\times p}\right)A_{\mu}(x)\star A_{\nu}(x')+\left(e^{-(\partial+\partial')\times p}-e^{(\partial-\partial')\times p}\right)A_{\nu}(x')\star A_{\mu}(x)\right]_{x'=x}\end{aligned} \tag{26}$$

Rewriting the Eq.(26) in the momentum space we find

$$\tilde{\Gamma}^{(2)} \;=\; \frac{i}{2}\frac{g_{\mathrm{M}}^2}{g_{\mathrm{D}}^2}\int\frac{\mathrm{d}^3k}{(2\pi)^3}\left(\tilde{\Pi}^{\mu\nu}_{\mathrm{p}}+\tilde{\Pi}^{\mu\nu}_{\mathrm{np}}\right)A_{\mu}(k)A_{\nu}(-k), \tag{27}$$

where the contributions from planar graphics are given by $\tilde{\Pi}^{\mu\nu}_{\mathrm{p}} = \Pi^{\mu\nu}$, and the contributions from non-planar graphics are written as

$$\tilde{\Pi}^{\mu\nu}_{\mathrm{np}} = -\frac{1}{2}\,\mathrm{tr}\int\frac{\mathrm{d}^3p}{(2\pi)^3}\left(S(p)\gamma^{\mu}S(p)\gamma^{\nu}+S(p)\gamma^{\mu}S(p)\not{k}S(p)\gamma^{\nu}+\cdots\right)\left(e^{2ik\times p}+e^{-2ik\times p}\right), \tag{28}$$

where we have used the Eq.(16). Thus, keeping only linear term in the derivative, the Eq.(28) turns out to be

$$\tilde{\Pi}^{\mu\nu}_{\mathrm{np}} \;=\; -m\,\mathrm{tr}\int\frac{\mathrm{d}^3p}{(2\pi)^3}\frac{e^{i\bar{\theta}.p}}{(p^2-m^2)^3}\times\left(\not{p}\gamma^{\mu}\not{p}\not{k}\gamma^{\nu}+\not{p}\gamma^{\mu}\not{k}\not{p}\gamma^{\nu}+\gamma^{\mu}\not{p}\not{k}\not{p}\gamma^{\nu}+m^2\gamma^{\mu}\not{k}\gamma^{\nu}\right), \tag{29}$$

with $\bar{\theta}^{\alpha} = \theta^{\alpha\beta}k_{\beta}$. Making use of the identities $\gamma^{\alpha}\gamma_{\alpha} = 3$,

$$\mathrm{tr}(\gamma^{\mu}\gamma^{\nu}\gamma^{\alpha}\gamma^{\beta}\gamma^{\rho}) = g^{\mu\nu}\mathrm{tr}(\gamma^{\alpha}\gamma^{\beta}\gamma^{\rho})+g^{\alpha\beta}\mathrm{tr}(\gamma^{\mu}\gamma^{\nu}\gamma^{\rho})+g^{\mu\rho}\mathrm{tr}(\gamma^{\alpha}\gamma^{\beta}\gamma^{\nu})-g^{\nu\rho}\mathrm{tr}(\gamma^{\alpha}\gamma^{\beta}\gamma^{\mu})$$

and of the relations for the integrals [30]

$$\begin{aligned}\int\frac{\mathrm{d}^3p}{(2\pi)^3}\frac{e^{i\bar{\theta}.p}}{(p^2-m^2)^3} \;&=\; -\frac{i}{32\pi}\frac{1}{m^2|m|}\left(1+m|\bar{\theta}|\right)e^{-m|\bar{\theta}|}\\ \int\frac{\mathrm{d}^3p}{(2\pi)^3}\frac{p_{\alpha}p_{\beta}\,e^{i\bar{\theta}.p}}{(p^2-m^2)^3} \;&=\; \frac{i}{32\pi}\frac{1}{|m|}\left(g_{\alpha\beta}-m|\bar{\theta}|\frac{\bar{\theta}_{\alpha}\bar{\theta}_{\beta}}{\bar{\theta}^2}\right)e^{-m|\bar{\theta}|},\end{aligned} \tag{30}$$

we can express the Eq.(29) in the form

$$\tilde{\Pi}^{\mu\nu}_{\mathrm{np}} \;=\; -\frac{k_{\rho}}{4\pi}\frac{m}{|m|}e^{-m|\bar{\theta}|}\epsilon^{\mu\nu\rho}-\frac{k_{\rho}}{16\pi}\frac{m^2}{|m|}|\bar{\theta}|e^{-m|\bar{\theta}|}\left[\epsilon^{\mu\nu\rho}-\frac{\bar{\theta}_{\alpha}}{\bar{\theta}^2}\left(\bar{\theta}^{\mu}\epsilon^{\alpha\nu\rho}+\bar{\theta}^{\nu}\epsilon^{\mu\alpha\rho}+\bar{\theta}^{\rho}\epsilon^{\mu\nu\alpha}\right)\right] \tag{31}$$

The second term of this equation disappears when we use the relation

$$\bar{\theta}_{\alpha}\left(\bar{\theta}^{\mu}\epsilon^{\alpha\nu\rho}+\bar{\theta}^{\nu}\epsilon^{\mu\alpha\rho}+\bar{\theta}^{\rho}\epsilon^{\mu\nu\alpha}\right) = \bar{\theta}^2\epsilon^{\mu\nu\rho}, \tag{32}$$

so that the non-planar graphics contribute as

$$\tilde{\Pi}^{\mu\nu}_{\rm np} = -\frac{k_\rho}{4\pi}\frac{m}{|m|}e^{-m|\bar{\theta}|}\epsilon^{\mu\nu\rho}. \tag{33}$$

Now substituting the expressions obtained for planar and non-planar graphics into Eq.(26), we obtain the term

$$\tilde{\Gamma}^{(2)} \; = \; -\frac{1}{8\pi}\frac{m}{|m|}\frac{g^2_{\rm M}}{g^2_{\rm D}}\int {\rm d}^3x\left(1-e^{-m|\bar{\theta}|}\right)\epsilon^{\mu\nu\rho}\partial_\mu A_\nu \star A_\rho. \tag{34}$$

The corresponding action for the cubic term, in the momentum space, is given by

$$\tilde{\Gamma}^{(3)} = \frac{i}{3}\frac{g^3_{\rm M}}{g^2_{\rm D}}\int\frac{{\rm d}^3k}{(2\pi)^3}\int\frac{{\rm d}^3k'}{(2\pi)^3}\left(\tilde{\Gamma}^{\mu\rho\nu}_{\rm p}+\tilde{\Gamma}^{\mu\rho\nu}_{\rm np}\right)e^{-ik'\times k}A_\mu(k')A_\rho(k)A_\nu(-k-k'). \tag{35}$$

The planar diagrams contribute as $\tilde{\Gamma}^{\mu\rho\nu}_{\rm p} = \Gamma^{\mu\rho\nu}$ and the expression for non-planar diagrams is given by

$$\tilde{\Gamma}^{\mu\rho\nu}_{\rm np} = -\frac{1}{2}\,{\rm tr}\int\frac{{\rm d}^3p}{(2\pi)^3}S(p)\gamma^\mu S(p)\gamma^\rho S(p)\gamma^\nu\left(e^{ik''\times p}+e^{-ik''\times p}\right), \tag{36}$$

where $k'' = -k - k'$. After manipulating this equation we can see that it is very similar to the Eq.(29) and therefore the Eq.(35) takes the form

$$\tilde{\Gamma}^{(3)} = \frac{ig_{\rm M}}{12\pi}\frac{m}{|m|}\frac{g^2_{\rm M}}{g^2_{\rm D}}\int {\rm d}^3x\left(1-e^{-m|\bar{\theta}|}\right)\epsilon^{\mu\nu\rho}A_\mu\star A_\nu\star A_\rho. \tag{37}$$

Finally, we combine the planar and non-planar contributions to get the Chern-Simons term for the Majorana representation, i.e.,

$$\tilde{\Gamma}_{\rm cs} = -\frac{1}{8\pi}\frac{m}{|m|}\frac{g^2_{\rm M}}{g^2_{\rm D}}\int {\rm d}^3x\left(1-e^{-m|\bar{\theta}|}\right)\epsilon^{\mu\nu\rho}\left(\partial_\rho A_\mu\star A_\nu - \frac{2ig_{\rm M}}{3}A_\mu\star A_\nu\star A_\rho\right). \tag{38}$$

In this equation we observe that the Chern-Simons action $\tilde{\Gamma}_{\rm cs}$ is vanish when we take $\bar{\theta}\to 0$. This means that UV/IR mixing, which is a characteristic of some noncommutative field theories, is not observed. Also, as we can see in the Eq.(38) if $m\to 0$, massless fermions, we do not have induced Chern-Simons action. This happens due to fact that the term $m\bar{\psi}\psi$ is vanish and therefore the parity symmetry is not broken. As in the commutative case parity violation is prerequisite to generate Chern-Simons term. We can still consider $m|\bar{\theta}|\to$ fixed, such that for $m\ll 1$ one has $|\bar{\theta}|\gg 1$. In this limit the Eq.(38) does not vanish and we predict light fermions traveling in a strongly noncommutative spacetime. Alternatively, from the dispersion relation, $E^2 = c^2p^2 + c^4m^2 + f(p^\mu\bar{\theta}_\mu)$, where $\bar{\theta}_\mu = \theta_{\mu\nu}p^\nu$, p_μ

here is the external momentum and f depends on the spin and charges of the particle — see, for instance, [31] and references therein — we see that the effects of the tensor $\theta_{\mu\nu}$ for light fermions, i.e., $m \ll 1$, with the same spin and charges, are stronger than the effects for heavy fermions. The effects of $\theta_{\mu\nu}$ on such particles could be, for instance, similar to the ones that a birefringent crystal induces in a light beam [31].

We summarize our work recalling that we have investigated the induction of a Chern-Simons action by the massive Thirring model in noncommutative spacetime, in the Dirac and Majorana representations. In the Dirac representation the Chern-Simons action is similar to ordinary non-abelian case.

As expected, the induced Chern-Simons action is gauge invariant under large gauge transformations. This happens due to the fact that we have considered parity violation that was originated in the fermion mass term. Another point that we have observed is that in the Majorana representation when we take $\bar{\theta} \to 0$ the Chern-Simons action is vanish and therefore UV/IR mixing does not appear.

Finally, we observe according to Eq.(38), by fixing $m|\bar{\theta}|$, we can predict that light fermions are more sensitive to the spacetime noncommutativity than heavy fermions.

Acknowledgments

We would like to thank D. Bazeia and V. O. Rivelles for useful discussions and CNPq, PROCAD for partial support.

[1] M. R. Douglas and N. A. Nekrasov, Rev. Mod. Phys.**73**, 977 (2001).

[2] R. J. Szabo, Quantum Field Theory on Noncommutative Space, hep-th/0109162.

[3] A. Connes, M.R. Douglas and A. Schwarz, J. High Energy Phys. **02**, 003 (1998).

[4] M. R. Douglas and C. Hull, J. High Energy Phys. **02**, 008 (1998).

[5] N. Seiberg and E. Witten, J. High Energy Phys. **09**, 032 (1999).

[6] R. Jackiw, Physical instances of noncommuting coordinates, hep-th/0110057.

[7] J. Gamboa, M. Loewe, F. Mendez, and J. C. Rojas, Mod.Phys.Lett.A **16**,2075 (2001).

[8] R. Prange and S. Girvin, The Quantum Hall effect Springer, New York, (1987).

[9] L. Susskind, The Quantum Hall Fluid and Noncommutative Chern-Simons Theory, hep-th/0101029.

[10] S. Deser, R. Jackiw, and S. Templeton, Ann. Phys. (N.Y.) **140**, 372 (1982).

[11] A. N. Redlich, Phys. Rev. Lett, **52**, 18, (1984); Phys. Rev. D **29**, 2366, (1984).

[12] M. Gomes, V. O. Rivelles and A. J. da Silva, Phys. Rev. D **41**, 1363 (1990).

[13] M. Gomes, R. S. Mendes, R. F. Ribeiro and A. J. da Silva, Phys. Rev. D **43**, 3516 (1991).

[14] T. Krajewski, Gauge invariance of the Chern-Simons action in noncommutative geometry, math-ph/9810015.

[15] A. A. Bichl, J. M. Grimstrup, V. Putz and M. Schweda, J. High Energy Phys. **07**, 046 (2000) [hep-th/0004071].

[16] G.-H. Chen and Y.-S. Wu, Nucl. Phys. **B593**, 562 (2001) [hep-th/0006114].

[17] C.-H. Chu, Nucl. Phys. **B580**, 352 (2000) [hep-th/0003007].

[18] N. Grandi and G.A. Silva, Phys. Lett. **B507**, 345 (2001) [hep-th/0010113].

[19] A. P. Polychronakos, J. High Energy Phys. **11**, 008 (2000) [hep-th/0010264].

[20] D. Bak, S. K. Kim, K.-S. Soh and J. H. Yee, Phys. Rev. D **64**, 025018 (2001) [hep-th/0102137].

[21] M.M. Sheikh-Jabbari, Phys. Lett. **B510**, 247 (2001) [hep-th/0102092].

[22] D. Bak, K.-M. Lee, J.-H. Park, Phys. Rev. Lett **87**, 030402 (2001) [hep-th/0102188].

[23] V. P. Nair and A. P. Polychronakos, Phys. Rev. Lett. **87**, 030403 (2001) [hep-th/0102181].

[24] S. Ghosh, Bosonization in the Noncommutative Plane, [hep-th/0303022].

[25] I. J. R. Aitchison and C. M. Fraser, Phys. Lett. **B146**, 63 (1984).

[26] A. I. Vanshtein, V. I. Zakharov, V. A. Novikov and M. A. Shifman, Yad. Fiz. **39** 124 (1984) [Sov. J. Nucl. Phys. **39**, 77 (1984)].

[27] C. M. Fraser, Z. Phys, C **28**, 101 (1985).

[28] I. J. R. Aitchison and C. M. Fraser, Phys. Rev. D **31** (1985) 2605.

[29] P. T. Matthews and A. Salam, Nuovo Cimento **12**, 563 (1954).

[30] F. T. Brandt, Ashok Das and J. Frenkel, Phys. Rev. D **65**, 085017 (2002) [hep-th/0112127].

[31] G. Amelino-Camelia, On the fate of Lorentz symmetry in loop quantum gravity and noncommutative spacetimes, [gr-qc/0205125].